

RESEARCH

Open Access



Dual harm among patients attending a mental health unit in Uganda: a hospital based retrospective study

Alain Favina¹, Joan Abaatyo¹ and Mark Mohan Kaggwa^{1,2*}

Abstract

Background Dual harm encompasses the complex interplay of the co-occurrence of self-harm and aggression. Individuals with dual harm may display a more hazardous pattern of harmful behaviors like homicide-suicide compared to people with sole harm. This study aimed to examine the presence of dual harm among general psychiatry inpatients in a mental health unit in Uganda.

Methods A retrospective chart review of 3098 inpatients from January 2018 to December 2021. Dual harm reported experience at admission was based on experiences of self-harm with harm to people or property or both. Logistic regression assessed the association between dual harm and sociodemographics and clinical characteristics.

Results A total of 29 (1%) patients experienced dual harm, with five having experienced self-harm with both harm to others and property, 23 with harm to people, and one with harm to property. Dual harm was statistically significantly associated with the male gender at bivariate analysis. However, there were no statistically significant factors associated with dual harm at multivariate analysis or sensitivity analysis with the specific types of dual harm.

Conclusion General psychiatry inpatients in Uganda experience dual harm before admission at lower prevalence than in previous literature. However, no investigated sociodemographic and clinical factors could explain these experiences. Further studies looking at dual harm are warranted to understand these unfortunate experiences with serious consequences among patients in Uganda.

Keywords Dual harm, Self-harm, Aggression, Male, Mental health, Uganda

Introduction

Dual harm encompasses the complex interplay of the co-occurrence of self-harm and aggression [1]. In addition to violent crimes, aggression can take many forms, such as verbal, psychological, physical, or toward other people or property [2]. On the other hand self-harm can take the form of suicidality or (non-suicidal) self-injury (NSSI) [1]. Individuals suffering from mental disorders often exhibit behaviors that pose risks to both themselves and the people or objects around them [3–5]. These people may have severe emotional turmoil, excessive rage, or a deep sense of hopelessness, which may manifest as violent outbursts

*Correspondence:

Mark Mohan Kaggwa
kmarkmohan@gmail.com

¹Department of Psychiatry, Faculty of Medicine, Mbarara University of Science and Technology, Mbarara, Uganda

²Department of Psychiatry and Behavioral Neurosciences, McMaster University, Hamilton, Canada



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

or destructive behavior toward other people or property [2, 6]. Additionally, these individuals may turn their aggression inwards, resorting to self-harming behaviors as a way to cope with their emotional pain [2, 7]. Contrasted to those who engage in self-harm alone or aggression alone (“sole-harm”), individuals who dual-harm may display a more hazardous pattern of harmful behaviors like homicide-suicide [1, 8–10].

A systematic review by Shafti et al., (2023) estimated that the overall prevalence of dual harm is above 20%, based on studies involving 15,094 individuals from different countries [10]. Previous research have also found that dual harm was more common among institutionalized individuals, especially male patients admitted to psychiatric hospitals [1, 11, 12] or correctional facilities [13–18]. However, dual harm was not limited to these populations, as it was also reported among patients in general psychiatry settings, university students, children and adolescents, homeless individuals, among others [19]. Despite extensive research on dual harm in psychiatry, there is a gap in the literature regarding the burden of dual harm among individuals in Africa.

The phenomenon of experiencing dual harm is associated with numerous multifaceted factors, which can be categorized into three domains: childhood experiences, biological factors, psychological factors and psychiatric diagnosis. (i) Childhood experiences: The family environment in which a person grows up can have a lasting impact on their propensity for dual harm. Studies have shown that negative childhood experiences within the family, such as abuse, household dysfunction, harsh discipline, and early neglect, are associated with aggression and dual harm [20–22]. Other family-related risk factors for dual harm include, poverty, family criminality, and educational underachievement [1, 23, 24]. Moreover, adverse childhood events, such as child abuse and parental death, increase the likelihood of dual harm [16]. (ii) Biological factors: Some biological factors, such as younger age and male sex, are related to dual harm [1, 2, 11]. Additionally, a person’s personality style may predispose them to harmful behaviors when biological elements and unfavorable environmental conditions coincide [9]. (iii) Psychological factors and mental health conditions: A person’s mental state and coping skills can also affect their tendency for dual harm. Individuals who experience depression or other mental health issues, use drugs or alcohol, observe self-harm, or are victims of violence are more likely to transition from sole harm at an early age to dual harm later in life [2]. Among adolescents, dual harm is linked to harsh parenting style, substance use, low school engagement, and low self-control [25, 26]. Furthermore, adolescents with high levels of anxiety and depressive symptoms are more prone to engage in dual harm [25]. Substance abuse disorders

and personality disorders (anti-social, borderline, and schizotypal) are significantly associated with dual harm in psychiatric patients [1]. Mood disorders are also documented to increase the risk of dual harm [16]. Additionally, adolescents who develop psychosis before the age of 18 years are more likely to engage in dual harm [25]. Trait impulsivity also predicts dual harm, especially in individuals with borderline personality disorder [27].

Dual harm poses a significant risk to individuals’ lives and well-being [9]. It has public health implications, as it contributes to increased healthcare utilization, emergency department visits, and psychiatric hospitalizations [20, 22, 28]. Additionally, it has profound social and economic costs, including the burden on healthcare systems, loss of productivity, and societal repercussions like the perpetuation of stigma surrounding mental health, community safety concerns and negative impact on family relationships [9, 20]. Moreover, it poses harm to both self and others, and its associated complications, such as psychological trauma among the victims, suicide, homicide, and pathological bereavement among those who have lost loved ones, may result in further burden to an already fragile mental health system [19]. Therefore, knowledge about factors associated with dual harm enables mental health professionals to develop targeted and effective treatment plans [29]. It facilitates the identification of individuals at higher risk, allowing for tailored interventions and preventive strategies and contributes to breaking down the stigma surrounding mental health issues, fostering a more compassionate and informed societal response [2, 9, 29]. Additionally, understanding the factors influencing dual harm contributes to the development of targeted and evidence-based treatments [29]. This can lead to improved outcomes for individuals experiencing mental health conditions associated with dual harm [29].

Adverse environmental factors like socio-economic disparities, political instability, and limited access to mental health services in Uganda [22, 30] may increase the likelihood of dual harm, rendering it a pressing concern. Moreover, studies examining dual harm in psychiatry in-patient settings, where the expectation that dual harm would be far more prevalent, are scarce [1, 12]. Therefore, the current study examined dual harm prevalence and associated sociodemographic and clinical factors among in-patient individuals in a mental health unit in Uganda.

Methods

This study was based on a retrospective chart review of the inpatient registered in Health Management Information Systems [HMIS] form 031 of the Mbarara Regional Referral Hospital (MRRH) psychiatry ward from January 2018 to December 2021. Similar data has been used to

understand various aspects of Uganda patients [31–35]. The current study was conducted in accordance with the Declaration of Helsinki 2013 and received a formal consent waiver from the Mbarara University Ethical Review Board, reference number MUST-2021-229.

MRRH psychiatry ward is the largest psychiatric facility in southwestern Uganda, with four stationed psychiatrists, and it's a teaching center for psychiatry residents, medical students, and nursing students. The diagnoses for mental illness are based on both ICD and or DSM criteria, following historical and observational content from the patient, family, and staff. The facility handles and manages people, especially referrals from all over southwestern Uganda. Information about the patients who are managed at MRRH is summarized in the HMIS. All patients in the HMIS have their records stored at the Department of Psychiatry records offices, and the records are managed by professional records managers. The HMIS form 031 captures the following information: date of admission, patient's number, name, address, age, anthropometric measurements, education level, occupation, gender, next of kin, investigations, presenting complaints, physical symptoms, diagnosis, comorbidities, and treatment given. Information from this form is used to physically trace the patient's chart, where the current data was extracted.

After retrieving all patient charts between January 2018 and December 2021 ($n=3014$), the research team created a database containing sociodemographic and clinical characteristics. However, the following variables were used to answer the current research questions: (i) patient's number, (ii) year of admission, (iii) age, (iv) gender (male, female), (v) marital status, (vi) level of education, (vii) employment status, (viii) presenting complaint including harm to self, others, and property, (ix) history of mental illness in the family, and (x) diagnosis. All data were entered parallel by two pairs of individuals. In case of any discrepancies, AF resolved them.

Data analysis

Data were analyzed using STATA version 17.0. Categorical variables were presented with frequencies and percentages. The continuous variable (age) was present with a mean and SD. The normality assumption was assessed using the Shapiro-Wilks test. The Chi-square tests for categorical variables were performed to determine significant differences between individuals with different types of aggression. The Fisher Exact test was performed to determine significant differences between individuals with different types of dual harm. The difference between the ages was assessed using the student's t-test. Bivariate and multiple logistic regressions were performed to assess the association between harm to others and harm to self and to assess the association between dual harm

and social demographics. The significant level was less than 5% for a 95% confidence interval.

Results

Patient's characteristics

A total of 3098 participants were included in this study, and slightly more than half of the participants were male (55.97%). According to marital status, married people were the most represented (45.95%), followed by single (40.90%). Participants with no formal or primary level of education represented 60.78%. The most represented diagnosis was bipolar disorder (37.67%), followed by Schizophrenia and other primary psychosis (33.36%), substance use disorder (12.54%), psychosis secondary to a general medical condition (8.67%), and lastly depression (7.75%). Most participants (62.07%) spent only one week in the ward during their admission.

Prevalence of aggressive behaviors and dual harm

A total of 29 (1%) individuals experienced dual harm. Five of these had dual harm to both people and property; 23 had dual harm to both others and self; and one individual had dual harm to self and property (see Fig. 1). The individual who had dual harm to self and property was a 49-year old divorced male farmer with a negative family history of mental illness.

Relationship between individual aggressive behaviors type and sociodemographic characteristics

Harm to people

Harm to people was statistically more among individuals diagnosed with bipolar disorder (p -value <0.001). On average, those who caused harm to people were younger than those who did not, i.e., mean of 32.27 vs. 34.92, p -value <0.001 . More males, 61.53%, caused harm to people than females, 38.47%, p -value <0.001 . Harm to people was statistically more among individuals who were single (p -value=0.009) and among those who had been on the ward for one week or less compared to those who had been admitted for more than a week (58.51% vs. 41.49, p -value=0.001) (Table 1).

Harm to property

Harm to property was statistically more among individuals diagnosed with bipolar disorder (p -value <0.001) and those who were single (p -value=0.001). On average, those who caused harm to property were younger than those who did not, i.e., mean of 31.94 vs. 34.41, respectively, p -value <0.001 . More males, 65.87%, caused harm to property than females, 34.13%, p -value <0.001 . More individuals who had been on the ward for one week (56.41%) caused harm to property than those who had been admitted for more than a week (43.59), p -value=0.006. Harm to property was statistically less



Fig. 1 Venn diagram showing the relation between the different aspects of aggression

among individuals with a positive family history of mental illness than those with a negative family history (22.69% vs. 72.31%, p -value=0.013) (Table 2).

Harm to self

Harm to self was statistically more among individuals diagnosed with a diagnosis of depression (p -value<0.001) (Table 3).

Factors associated with dual harm

Bivariate and multivariate analysis was done to identify factors associated with dual harm. At bivariate analysis, only male sex was associated with dual harm (odds ratio (OR): 02.49; 95% confidence interval (CI): 01.06–05.85; p -value=0.036). However, there was no factor associated

with dual harm at multivariate analysis (Table 4). Even at sensitivity analysis, no factors were identified to be associated with the specific types of dual harm.

Discussion

This study aimed to investigate dual harm and its related factors among individuals admitted with mental illness. Results showed that the prevalence of dual-harm among these individuals admitted at MRRH was approximately 1%, which is closer to that of 2% reported from hospital records of general psychiatry inpatients in USA [5]. The prevalence reported in this study is much lower than that of 17% reported among general psychiatry outpatients in UK [36], and that of 20% among inpatients in Netherlands [7], and in Finland [37]. The higher prevalence is

Table 1 Relationship between harm to people and sociodemographic characteristics

Variables	Harm To people		
	No	Yes: n = 1338 (43.19%)	P value
Age (mean, SD)	34.92, 14.69	32.27, 12.93)	< 0.001
Sex			
Female	848, 48.24%	514, 38.47%	< 0.001
Male	910, 51.76%	822, 61.53%	
Marital status			
Divorced/Separated/widowed	191, 13.40%	139, 12.78%	0.009
Single	546, 38.32%	482, 44.30%	
Married/cohabiting	688, 48.28%	467, 42.92%	
Level of education			
None	374, 31.30%	266, 29.62%	0.309
Primary	344, 28.79%	288, 32.07%	
Secondary	282, 23.60%	215, 23.94%	
Tertiary	195, 16.32%	129, 14.37%	
Occupation			
Unemployed	235, 17.07%	175, 16.43%	0.816
Pupil/Student	198, 14.38%	159, 14.93%	
Farmer	442, 32.10%	357, 33.52%	
Other employment	502, 36.46%	374, 35.12%	
Diagnosis			
Bipolar	414, 32.50%	450, 44.12%	< 0.001
Schizophrenia	434, 34.07%	331, 32.45%	
Depression	158, 12.40%	20, 1.96%	
Psychosis secondary to GMC*	128, 10.05%	71, 6.96%	
Substance use disorder	140, 10.99%	148, 14.51%	
Duration on ward			
One week	1,061, 64.73%	739, 58.51%	0.001
Over one week	578, 35.27%	524, 41.49%	
Family History of mental illness			
positive	339, 19.53%	234, 17.66%	0.189
Negative	1,397, 80.47%	1,091, 82.34%	

*General medical condition

Table 2 Relationship between harm to property and sociodemographic characteristics

Variables	Harm To Property		
	No	Yes: n = 502 (16.20%)	P value
Age	34.14, 14.08	31.94, 13.55	0.001
Sex			
Female	1,191, 45.93%	171, 34.13%	< 0.001
Male	1,402, 54.07%	330, 65.87%	
Marital status			
Divorced/Separated/widowed	283, 13.55%	47, 11.08%	0.001
Single	821, 39.30%	207, 48.82%	
Married/cohabiting	985, 47.15%	170, 40.09%	
Level of education			
None	505, 29.74%	135, 34.18%	0.001
Primary	496, 29.21%	136, 34.43%	
Secondary	412, 24.26%	85, 21.52%	
Tertiary	285, 16.78%	39, 9.87%	
Occupation			
Unemployed	327, 16.10%	83, 20.19%	0.210
Pupil/Student	298, 14.67%	59, 14.36%	
Farmer	666, 32.79%	133, 32.36%	
Other employment	740, 36.44%	136, 33.09%	
Diagnosis			
Bipolar	689, 36.09%	175, 45.45%	< 0.001
Schizophrenia	635, 33.26%	130, 33.77%	
Depression	171, 8.96%	7, 1.82%	
Psychosis secondary to GMC*	181, 9.48%	18, 4.68%	
Substance use disorder	233, 12.21%	55, 14.29%	
Duration on ward			
One week	1,536, 63.11%	264, 56.41%	0.006
Over one week	898, 36.89%	204, 43.59%	
Family history of mental illness			
positive	460, 17.95%	113, 22.69%	0.013
Negative	2,103, 82.05%	385, 77.31%	

likely because the study in UK assessed for prior history of violence among those who had attempted suicide and the study in Netherlands used tools to assess for aggression and self-harm while this study denoted aggression reported at admission. Another possible reason for the

Table 3 Relationship between self-harm and sociodemographic characteristics

Variables	Harm To Self		P value
	No	Yes: n = 107 (3.45%)	
Age	33.83, 14.05	32.49, 13.06	0.331
Sex			
Female	1,319, 44.16%	43, 40.19%	0.430
Male	1,668, 55.84%	64, 59.81%	
Marital status			
Divorced/Separated/widowed	323, 13.36%	7, 7.37%	0.222
Single	988, 40.86%	40, 42.11%	
Married/cohabiting	107, 45.78%	48, 50.53%	
Level of education			
None	621, 30.74%	19, 26.03%	0.411
Primary	613, 30.35%	19, 26.03%	
Secondary	474, 23.47%	23, 31.51%	
Tertiary	312, 15.45%	12, 16.44%	
Occupation			
Unemployed	394, 16.74%	16, 18.18%	0.630
Pupil/Student	341, 14.49%	16, 18.18%	
Farmer	775, 32.92%	24, 27.27%	
Other employment	844, 35.85%	32, 36.36%	
Diagnosis			
Bipolar	852, 38.41%	12, 15.79%	<0.001
Schizophrenia	745, 33.59%	20, 26.32%	
Depression	146, 6.58%	32, 42.11%	
Psychosis secondary to GMC*	198, 8.93%	1, 1.32%	
Substance use disorder	277, 12.49%	11, 14.47%	
Duration on ward			
One week	1,736, 62.00%	64, 62.75%	0.879
Over one week	1,064, 38.00%	38, 37.25%	
Family History of mental illness			
positive	551, 18.65%	22, 20.56%	0.614
Negative	2,403, 81.35%	85, 79.44%	

higher prevalence in the study in Finland is that it was carried out for a longer period (16years) and it focused on homicide-suicide dual harm while the current study looked at non-homicide-suicide dual harm. The prevalence in this study is also much lower than that reported among forensic psychiatry populations [10, 38, 39] likely because forensic populations often consist of individuals who may exhibit complex psychopathological conditions, which can involve a combination of traits, such as impulsivity, emotional dysregulation, and interpersonal difficulties [40]. Additionally, cultural norms and values in Uganda may prioritize communal and family support more [41], potentially resulting in lower rates of dual harm. The robust family networks in Uganda may offer greater care and supervision to individuals with mental disorders [42–44], thus reducing the likelihood of dual harm. In addition, cultural norms in Uganda play a significant role in shaping the behavior and help-seeking patterns of psychiatry patients, often discouraging the presentation of violence in healthcare settings. Family members of violent people with mental illness may be reluctant to seek help at hospitals or clinics because of Africa's societal norms favoring restraint, which includes physically restraint of the mentally ill, and social cohesion [45, 46]. This is because they may be more worried about maintaining their social connections and reputation [46]. There may also be reporting bias as a result of these cultural norms, when people believe they shouldn't expose their patients' violent occurrences to the public since doing so would make them appear terrible [47, 48]. Lastly, aggression is a multifaceted behavior that encompasses various forms, and it remains unclear whether the inpatient charts of psychiatry patients comprehensively captured all forms of aggression like verbal aggression. However, the culture of Ugandan people is changing [49] and the trends may also increase with time.

This study found that being male was significantly associated with dual harm. This finding echoes findings in other studies [11, 50]. This is likely because males may be more likely to engage in risky or dangerous behaviors potentially leading to dual harm [51]. Males resort to aggressive methods in order to ascertain their masculinity, potentially leading to their involvement in dual harm [52, 53]. Additionally, there are higher rates of substance abuse among men, such as alcohol or illicit drugs, can increase the likelihood of aggressive or violent behavior, both toward oneself and others [54–56]. Men might have smaller social support networks or feel less comfortable discussing their mental health issues with others [57], which can lead to increased isolation and a higher risk of harm. Despite the identified relationship at bivariate regression, it was not significant in the multivariate model, an indication of influence of other factors in its

Table 4 Factors associated with the dual harm

	Dual harm			Bivariate analysis			Multivariate analysis		
	Absent	Present	χ^2 (p value)	OR	95% CI	p-value	AOR	95% CI	p-value
	<i>n=29</i>								
Age	33.82 (14.02)	29.59 (13.48)	(t value missing) 0.105	00.97	(00.94–01.01)	0.107	00.95	(00.88–01.03)	0.224
Sex									
Female	1,355 44.21%	7 24.14%	4.70 (0.370)	01.00			01.00		
Male	1,710 55.79%	22 75.86%		02.49	(01.06–05.85)	0.036	01.49	(00.45–04.94)	0.514
Marital status									
Divorced/Separated/widowed	329 13.23%	1 3.70%	3.44 (0.199)	01.00			01.00		
Single	1,013 40.75%	15 55.56%		04.87	(00.64–37.02)	0.126	01.18	(00.28–04.99)	0.824
Married/cohabiting	1,144 46.02%	11 40.74%		03.16	(00.41–24.59)	0.271	01.00	Omitted	
Level of education									
None	637 30.74%	3 14.29%	2.66 (0.375)	01.00			01.00		
Primary	624 30.12%	8 38.10%		02.72	(00.72–10.31)	0.140	02.15	(00.41–11.40)	0.367
Secondary	491 23.70%	6 28.57%		02.59	(00.65–10.43)	0.179	02.23	(00.38–12.98)	0.373
Tertiary	320 15.44%	4 19.05%		02.65	(00.59–11.93)	0.203	02.24	(00.34–14.92)	0.404
Occupation									
Unemployed	403 16.66%	7 30.43%	3.12 (0.411)	01.00			01.00		
Pupil/Student	354 14.63%	3 13.04%		00.49	(00.13–01.90)	0.301	00.27	(00.05–01.29)	0.101
Farmer	793 32.78%	6 26.09%		00.44	(00.15–01.30)	0.138	00.15	(00.02–01.31)	0.085
Other employment	869 35.92%	7 30.43%		00.46	(00.16–01.33)	0.153	00.55	(00.15–02.04)	0.373
Diagnosis									
Substance use disorder	283 12.47%	5 20.83%	3.06 (0.505)	01.00			01.00		
Schizophrenia	757 33.35%	8 33.33%		00.60	(00.19–01.84)	0.371	00.50	(00.12–02.14)	0.352
Depression	175 7.71%	3 12.50%		00.97	(00.23–04.11)	0.967	00.67	(00.06–07.34)	0.746
Secondary	198 8.72%	1 4.17%		00.29	(00.03–02.47)	0.255	00.67	(00.06–07.37)	0.745
Bipolar	857 37.75%	7 29.17%		00.46	(00.15–01.47)	0.191	00.42	(00.09–01.85)	0.250
Weeks in ward									
one Week	1,786 62.14%	14 50.00%	1.74 (0.244)	01.00			01.00		
Over one week	1,088 37.86%	14 50.00%		01.64	(00.78–03.46)	0.192	01.91	(00.69–05.30)	0.215
History of stressors									
No	568 18.73%	5 17.24%	0.04 (1.00)	01.00			01.00		
Yes	2,464 81.27%	24 82.76%		01.11	(00.42–02.91)	0.838	01.70	(00.59–04.93)	0.328

relationship with dual harm. An aspect that may be due to the limitation of the present study.

In this study, no specific mental disorder was associated with dual harm. This finding aligns with finding of a systematic review from twenty-seven studies that there is no sufficient evidence that any of the mental illnesses is associated with dual-harm when compared [10]. This implies that various mental disorders are not unique to dual harm, but rather driven by the separate self-harm or aggressive behaviors [10].

Contrary to various studies where prior adverse events resulted in dual harm [23, 24], this study's findings do not report any association between history of a stressor and dual harm. This is likely because Uganda has a unique cultural context, and individuals may have different coping mechanisms and ways of expressing psychological distress. Additionally, social support [58] or individual resilience [59], could have influenced the outcomes. Furthermore, the way in which stressors were measured and reported in the study could introduce biases i.e., stressors were self-reported, and participants may underreport or fail to recognize certain stressors. In addition, the study might not have captured the delayed effects of long-term stressors.

The absence of identified factors associated with dual harm in the final model of this study can be attributed to the complexities and multifaceted nature of mental health outcomes. Several variables may influence the occurrence of dual harm, but they were difficult to capture comprehensively within the confines of this study as reported findings from history captured at admission. Furthermore, identifying predictors of dual harm might be difficult due to characteristics including individual resilience, distinct coping techniques, and subtle manifestations of discomfort among Ugandan patients. Moreover, the model's capacity to identify meaningful correlations may be impacted by methodological constraints like the duration taken into consideration and the absence of measuring instruments. The interaction of these variables, together with the variety of psychiatric disorders and personal experiences, highlights the need for more thorough and situation-specific study to gain a deeper understanding of the elements causing twofold harm among Ugandan psychiatric in-patients.

Overall, the study's finding that 1% of psychiatric in-patients in Uganda experience dual harm, has significant ramifications for mental health services and research in the area. Recognizing dual arm can help mental health providers create sensitive treatment plans that meet the difficulties the patients encounter. These plans may include strengthening coping skills and decreasing aggressive behavior. These results further highlight the importance of increasing mental health service

accessibility and increasing dual harm awareness in order to enhance early detection and intervention.

Limitations and recommendations

The following limitations should be taken into consideration when interpreting the results of this study. Firstly, it should be noted that this was a retrospective evaluation, and study designs like these frequently have missing data. In a study of this nature, the limitations of missing data are universally recognized. We recommend future researchers to use several assessment techniques and employ prospective study designs to ensure accurate detection of the various aspects of dual harm. Secondly, clinical judgment was used for assessment and categorization of the harm because the nature of the harm was documented by the doctors in the patient's charts. The reported prevalence of dual harm among psychiatry in-patients may not be accurate or consistent due to subjectivity reporting by the caregivers. In order to prevent this bias, future investigators can employ formal techniques and operational criteria for dual-harm aspects, i.e., violence and self-harm. Thirdly, many variables that could have influenced the findings, such as adherence to medication, adverse childhood events, family environment were not retrieved from the records of patients and hence not included in the analysis. We recommend that prospective study designs be used in subsequent research to obtain this kind of data. Furthermore, the study population is limited to in-patients of general psychiatry and excludes members of the forensic psychiatry population because the psychiatric unit is not gazetted to house forensic patents, who frequently face unique clinical and legal circumstances that put them at a higher risk of harm. Hence, the lack of this particular patient category in our research may restrict the generalization of our conclusions to a larger psychiatric community and may understate the general prevalence of dual harm among inpatient psychiatric patients.

Conclusion

Dual harm is not a common phenomenon documented among inpatients with mental illness at a tertiary facility in Uganda. According to our research findings, no factors were found to be statistically linked with dual harm among psychiatry in-patients. This indicates that dual harm in this community is complicated and varied, highlighting the need for more research and sophisticated methods to comprehend the variables influencing such occurrences. Dual harm is still less explored and further research is need in completely understand this phenomenon in Uganda and Africa at large.

Author contributions

AF– Data collection, data analysis, initial draft, conceptualization, validation.
MMK– Conceptualization, data curation, visualization, validation, initial

manuscript, subsequent revisions, supervision. JA– Data collection, drafting of the initial manuscript, subsequent revisions.

Funding

No funding source is associated with this work.

Data availability

The datasets will be made available to appropriate academic parties on request from the corresponding author.

Declarations

Ethics approval and consent to participate

The study was approved by Mbarara University of Science and Technology Research Ethics Committee (Approval number: MUST-2021-229) and provided a waiver of informed consent for study participants.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Received: 3 November 2023 / Accepted: 25 January 2024

Published online: 22 February 2024

References

- Slade K. Dual harm: the importance of recognising the duality of self-harm and violence in forensic populations. Volume 59. London, England: SAGE Publications Sage UK; 2019. pp. 75–7.
- Steed S, Farooq B, Taylor P, Shafti M, Mars B, Kapur N, Webb RT. Childhood predictors of self-harm, externalised violence and transitioning to dual harm in a cohort of adolescents and young adults. *Psychol Med* 2023;1–11.
- Edlinger M, Rauch AS, Kemmler G, Yalcin-Siedentopf N, Fleischhacker WW, Hofer A. Risk of violence of inpatients with severe mental illness—do patients with schizophrenia pose harm to others? *Psychiatry Res*. 2014;219(3):450–6.
- Sadath A, Troya MI, Nicholson S, Cully G, Leahy D, Ramos Costa AP, Benson R, Corcoran P, Griffin E, Phillip E, et al. Physical and mental illness comorbidity among individuals with frequent self-harm episodes: a mixed-methods study. *Front Psychiatry*. 2023;14:1121313.
- Tardiff K, Sweillam A. Assault, suicide, and Mental Illness. *Arch Gen Psychiatry*. 1980;37(2):164–9.
- Kulper DA, Kleiman EM, McCloskey MS, Berman ME, Coccaro EF. The experience of aggressive outbursts in intermittent explosive disorder. *Psychiatry Res*. 2015;225(3):710–5.
- Nijman HL, à Campo JM. Situational determinants of inpatient self-harm. *Suicide and Life-Threatening Behavior*. 2002;32(2):167–75.
- Steed S, Webb RT, Mok PL, Pedersen CB, Antonsen S, Kapur N, Carr MJ. Risk of dying unnaturally among people aged 15–35 years who have harmed themselves and inflicted violence on others: a national nested case-control study. *The Lancet Public Health*. 2019;4(5):e220–8.
- Shafti M, Taylor PJ, Forrester A, Pratt D. The co-occurrence of self-harm and aggression: a cognitive-emotional model of dual-harm. *Front Psychol*. 2021;12:586135.
- Shafti M, Taylor P, Forrester A, Handerer F, Pratt D. A systematic review of the co-occurrence of self-harm and aggression: is dual harm a unique behavioural construct? *Front Psychiatry*. 2023;14:1083271.
- Plutchik R, Van Praag HM, Conte HR. Correlates of suicide and violence risk: III. A two-stage model of countervailing forces. *Psychiatry Res*. 1989;28(2):215–25.
- Kottler C, Smith JG, Bartlett A. Patterns of violence and self-harm in women prisoners: characteristics, co-incidence and clinical significance. *J Forensic Psychiatr Psychol*. 2018;29(4):617–34.
- Slade K. Dual harm: an exploration of the presence and characteristics for dual violence and self-harm behaviour in prison. *J Criminal Psychol*. 2018;8(2):97–111.
- Mandelli L, Carli V, Roy A, Serretti A, Sarchiapone M. The influence of childhood trauma on the onset and repetition of suicidal behavior: an investigation in a high risk sample of male prisoners. *J Psychiatr Res*. 2011;45(6):742–7.
- Gunilla Stålenheim E. Relationships between attempted suicide, temperamental vulnerability, and violent criminality in a Swedish forensic psychiatric population. *Eur Psychiatry*. 2001;16(7):386–94.
- Huang Y, Zhang S, Zhong S, Gou N, Sun Q, Guo H, Lin R, Guo W, Chen H, Wang J, et al. The association of childhood adversities and mental health problems with dual-harm in individuals with serious aggressive behaviors. *BMC Psychiatry*. 2022;22(1):385.
- Slade K, Forrester A, Baguley T. Coexisting violence and self-harm: dual harm in an early-stage male prison population. *Legal and Criminological Psychology*. 2020;25(2):182–98.
- Maden A, Chamberlain S, Gunn J. Deliberate self-harm in sentenced male prisoners in England and Wales: some ethnic factors. *Criminal Behav Mental Health*. 2000;10(3):199–204.
- O'Donnell O, House A, Waterman M. The co-occurrence of aggression and self-harm: systematic literature review. *J Affect Disord*. 2015;175:325–50.
- Carr MJ, Steed S, Mok PLH, Pedersen CB, Antonsen S, Kapur N, Webb RT. Adverse childhood experiences and risk of subsequently engaging in self-harm and violence towards other people—dual harm. *Int J Environ Res Public Health* 2020, 17(24).
- Richmond-Rakerd LS, Caspi A, Arseneault L, Baldwin JR, Danese A, Houts RM, Matthews T, Wertz J, Moffitt TE. Adolescents who Self-Harm and commit violent crime: testing early-life predictors of dual harm in a Longitudinal Cohort Study. *Am J Psychiatry*. 2019;176(3):186–95.
- Steinhoff A, Bechtiger L, Ribeaud D, Eisner M, Shanahan L. Self-, other-, and dual-harm during adolescence: a prospective-longitudinal study of childhood risk factors and early adult correlates. *Psychol Med*. 2023;53(9):3995–4003.
- Flynn S, Gask L, Appleby L, Shaw J. Homicide–suicide and the role of mental disorder: a national consecutive case series. *Soc Psychiatry Psychiatr Epidemiol*. 2016;51:877–84.
- Rouchy E, Germanaud E, Garcia M, Michel G. Characteristics of homicide-suicide offenders: a systematic review. *Aggress Violent Beh*. 2020;55:101490.
- Slade K, Smith HP, Potter A, Baguley T. Re-examining the dual harm profile: an assessment using US prison population-level data. *Psychol Crime Law*:1–15.
- Spaan P, Michielsens PJS, de Neve-Enthoven NGM, Bouter DC, Grootendorst-van Mil NH, Hoogendijk WJG, Roza SJ. Dual-harm in adolescence and associated clinical and parenting factors. *Soc Psychiatry Psychiatr Epidemiol*. 2022;57(8):1615–26.
- Terzi L, Martino F, Berardi D, Bortolotti B, Sasdelli A, Menchetti M. Aggressive behavior and self-harm in Borderline personality disorder: the role of impulsivity and emotion dysregulation in a sample of outpatients. *Psychiatry Res*. 2017;249:321–6.
- Chung J, Sadeghzadeh K, Sibdari S. Psychiatric Hospitalization Associated with Emergency Department visits. *Issues Ment Health Nurs*. 2022;43(6):552–9.
- Ghossoub E, Cherro M, Akil C, Gharzeddine Y. Mental illness and the risk of self- and other-directed aggression: results from the National Survey on Drug Use and Health. *J Psychiatr Res*. 2021;132:161–6.
- Kaggwa MM, Harms S, Mamun MA. Mental health care in Uganda. *The Lancet Psychiatry*. 2022;9(10):766–7.
- Kaggwa MM, Abaatyo J, Alol E, Muwanguzi M, Najjuka SM, Favina A, Rukundo GZ, Ashaba S, Mamun MA. Substance use disorder among adolescents before and during the COVID-19 pandemic in Uganda: retrospective findings from a psychiatric ward registry. *PLoS ONE*. 2022;17(5):e0269044.
- Abaatyo J, Kaggwa MM, Favina A, Olagunju AT. Readmission and associated clinical factors among individuals admitted with bipolar affective disorder at a psychiatry facility in Uganda. *BMC Psychiatry*. 2023;23(1):474.
- Abaatyo J, Favina A, Bulega SE, Kaggwa MM. Suicidal behavior among inpatients with severe mental conditions in a public mental health hospital in Uganda. *BMC Psychiatry*. 2023;23(1):346.
- Abaatyo J, Favina A, Kaggwa MM. Absconding among admitted patients with bipolar affective disorder diagnosis in Uganda. *BMC Psychiatry*. 2023;23(1):318.
- Kaggwa MM, Rukundo GZ, Wakida EK, Maling S, Sserumaga BM, Atim LM, Obua C. Suicide and suicide attempts among patients attending primary Health Care facilities in Uganda: A Medical Records Review. *Risk Manage Healthc Policy*. 2022;15(null):703–11.
- HUNT IM, KAPUR N, WINDFUHR K, ROBINSON J, BICKLEY H, PARSONS FLYNN S, BURNS R, SHAW J. Suicide in Schizophrenia: findings from a National Clinical Survey. *J Psychiatric Practice*®. 2006;12(3):139–47.
- Virkkunen M. Suicide linked to homicide. *Psychiatr Q*. 1974;48(2):276–82.

38. Haggård-Grann U, Hallqvist J, Långström N, Möller J. Short-term effects of psychiatric symptoms and interpersonal stressors on criminal violence. *Soc Psychiatry Psychiatr Epidemiol*. 2006;41(7):532–40.
39. Boxer P. Covariation of self- and other-directed aggression among inpatient youth: continuity in the transition to treatment and shared risk factors. *Aggressive Behav*. 2010;36(3):205–17.
40. Garofalo C, Neumann CS, Velotti P. Difficulties in emotion regulation and psychopathic traits in violent offenders. *J Criminal Justice*. 2018;57:116–25.
41. Belaid L, Ochola E, Bayo P, Alii GW, Ogwang M, Greco D, Zarowsky C. Exploring the impact of a community participatory intervention on women's capability: a qualitative study in Gulu Northern Uganda. *BMC Womens Health*. 2021;21(1):28.
42. Olwit C, Musisi S, Leshabari S, Sanyu I. Chronic sorrow: lived experiences of caregivers of patients diagnosed with schizophrenia in Butabika Mental Hospital, Kampala, Uganda. *Arch Psychiatr Nurs*. 2015;29(1):43–8.
43. Kaggwa MM, Najjuka MS, Kesande C, Nyemara N, Kule M, Mamum MA, Bongomin F, Ashaba S. Length of stay of hospitalized patients at tertiary psychiatry facilities in Uganda: the role of caregiver's presence. *Discover Mental Health*. 2022;2(1):15.
44. Kaggwa MM, Najjuka SM, Mamun MA, Griffiths MD, Nyemara N, Ashaba S. Involvement and burden of informal caregivers of patients with mental illness: the mediating role of affiliated stigma. *BMC Psychiatry*. 2023;23(1):72.
45. Verity F, Turiho A, Mutamba BB, Cappo D. Family care for persons with severe mental illness: experiences and perspectives of caregivers in Uganda. *Int J Mental Health Syst*. 2021;15(1):48.
46. Asher L, Fekadu A, Teferra S, De Silva M, Pathare S, Hanlon C. I cry every day and night, I have my son tied in chains: physical restraint of people with schizophrenia in community settings in Ethiopia. *Globalization and Health*. 2017;13(1):47.
47. Annan J, Brier M, Aryemo F. From rebel to returnee daily life and reintegration for young soldiers in Northern Uganda. *J Adolesc Res*. 2009;24(6):639–67.
48. Orley JH. *Culture and mental illness: a study from Uganda*. East African Publishing House; 1970.
49. Sylvia N. The role of media in culture modification in Uganda: a case study of Toro Culture. 2007.
50. Dudeck M, Susic-Vasic Z, Otte S, Rasche K, Leichauer K, Tippelt S, Shenar R, Klingner S, Vasic N, Streb J. The association of adverse childhood experiences and appetitive aggression with suicide attempts and violent crimes in male forensic psychiatry inpatients. *Psychiatry Res*. 2016;240:352–7.
51. Melzer SA. Gender, work, and intimate violence: men's occupational violence spillover and compensatory violence. *J Marriage Family*. 2002;64(4):820–32.
52. Cawley J, Ruhm CJ. The economics of risky health behaviors. *Handbook of health economics*. Volume 2. edn.: Elsevier; 2011. pp. 95–199.
53. Kaggwa MM, Muwanguzi M, Nduhuura E, Kajjimu J, Arinaitwe I, Kule M, Najjuka SM, Rukundo GZ. Suicide among Ugandan university students: evidence from media reports for 2010–2020. *BJPsych Int*. 2021;18(3):63–7.
54. Chermack ST, Fuller BE, Blow FC. Predictors of expressed partner and non-partner violence among patients in substance abuse treatment. *Drug Alcohol Depend*. 2000;58(1–2):43–54.
55. Boles SM, Miotto K. Substance abuse and violence: a review of the literature. *Aggress Violent Beh*. 2003;8(2):155–74.
56. Atim LM, Kaggwa MM, Mamum MA, Kule M, Ashaba S, Maling S. Factors associated with elder abuse and neglect in rural Uganda: a cross-sectional study of community older adults attending an outpatient clinic. *PLoS ONE*. 2023;18(2):e0280826.
57. McKenzie SK, Collings S, Jenkin G, River J. Masculinity, Social Connectedness, and Mental Health: men's diverse patterns of practice. *Am J Men's Health*. 2018;12(5):1247–61.
58. Seffren V, Familiar I, Murray SM, Augustinavicius J, Boivin MJ, Nakasujja N, Opoka R, Bass J. Association between coping strategies, social support, and depression and anxiety symptoms among rural Ugandan women living with HIV/AIDS. *AIDS Care*. 2018;30(7):888–95.
59. Klasen F, Oettingen G, Daniels J, Post M, Hoyer C, Adam H. Posttraumatic resilience in former Ugandan child soldiers. *Child Dev*. 2010;81(4):1096–113.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.